

Frequently Asked Questions

What is a multiple pregnancy?

A pregnancy with more than one [fetus](#) is called a [multiple pregnancy](#).

How does multiple pregnancy happen?

If more than one [egg](#) is released during the [menstrual cycle](#) and each egg is fertilized by a sperm, more than one [embryo](#) may implant and grow in the [uterus](#). This type of pregnancy results in [fraternal twins](#) (or more). When a single fertilized egg splits, it results in multiple identical embryos. This type of pregnancy results in [identical twins](#) (or more). Fraternal twins are more common than identical twins.

What are some causes of multiple pregnancy?

The use of fertility drugs to induce [ovulation](#) often causes more than one egg to be released from the [ovaries](#) and can result in twins, triplets, or more.

[In vitro fertilization \(IVF\)](#) can lead to a multiple pregnancy if more than one embryo is transferred to the uterus. Identical multiples also may result if the fertilized egg splits after transfer.

Women older than age 35 are more likely to release two or more eggs during a single menstrual cycle than younger women. So older women are more likely than younger women to become pregnant with multiples.

What are some symptoms of multiple pregnancy?

Women who are pregnant with multiples may have more severe morning sickness or breast tenderness than women who are pregnant with a single fetus. They also may gain weight more quickly. Most multiple pregnancies are discovered during an [ultrasound exam](#).

Do I need to gain extra weight if I am pregnant with multiples?

Most women who are pregnant with multiples need to gain more weight than women who are pregnant with one fetus. An extra 300 calories a day is needed for each fetus. For example, if you are pregnant with twins, you need an extra 600 calories a day. For triplets and more, talk with your [obstetrician–gynecologist \(ob-gyn\)](#) about the right amount of weight gain for your pregnancy.

Should I exercise if I am pregnant with multiples?

Staying active during multiple pregnancy is important for your health, but you may need to avoid strenuous exercise. Try low-impact exercise, such as swimming, prenatal yoga, and walking. You should aim for 30 minutes of exercise a day. If problems arise during your pregnancy, your ob-gyn may recommend that you avoid exercise.

What will happen during my prenatal care?

You likely will have more frequent [prenatal care](#) visits with your ob-gyn. Starting in your second trimester, you may have ultrasound exams every 4 to 6 weeks. If a problem is suspected, you may have special tests or more frequent ultrasound exams.

What is the most common complication of multiple pregnancy?

The most common [complication](#) of multiple pregnancy is [preterm](#) birth. More than one half of all twins are born preterm. Triplets and more are almost always born preterm.

What are the risks of preterm birth?

Babies born before 37 weeks of pregnancy may have an increased risk of short-term and long-term health problems, including problems with breathing, eating, and staying warm. Other problems, such as learning and behavioral disabilities, may appear later in

childhood or even in adulthood. Very preterm babies (those who are born before 32 weeks of pregnancy) can die or have severe health problems, even with the best of care.

Preterm multiples also have a greater risk of serious complications that can lead to [cerebral palsy](#). Children born with problems related to being preterm may need lifelong medical care.

What are chorionicity and amnionicity?

Early in a multiple pregnancy, an ultrasound exam is done to find out whether each fetus has its own [chorion](#) ([chorionicity](#)) and [amniotic sac](#) ([amnionicity](#)). There are three types of twins:

1. Dichorionic–diamniotic—Twins who have their own chorions and amniotic sacs. They typically do not share a [placenta](#) and can be fraternal or identical.
2. Monochorionic–diamniotic—Twins who share a chorion but have separate amniotic sacs. They share a placenta and are identical.
3. Monochorionic–monoamniotic—Twins who share one chorion and one amniotic sac. They share a placenta and are identical.

What are the risks for with monochorionic twins?

Monochorionic twins have a higher risk of complications than those with separate placentas. One problem that can occur in monochorionic–diamniotic babies is [twin–twin transfusion syndrome \(TTTS\)](#). In TTTS, the blood flow between the twins becomes unbalanced. One twin donates blood to the other twin. The donor twin has too little blood, and the recipient twin has too much blood. The earlier TTTS occurs in the pregnancy, the more serious the outcomes for one or both babies.

Although monochorionic–monoamniotic babies are rare, this type of pregnancy is very risky. The most common problem is an [umbilical cord](#) complication. Women with this type of pregnancy are monitored more frequently and are likely to have a [cesarean birth](#).

How can multiple pregnancy affect my risk of preeclampsia?

Preeclampsia is a blood pressure disorder that usually starts after 20 weeks of pregnancy or after childbirth. It occurs more often in multiple pregnancies than in singleton pregnancies. It also tends to occur earlier and is more severe in multiple pregnancies.

Preeclampsia can damage many organs in your body, most commonly your kidneys, liver, brain, and eyes. Preeclampsia that worsens and causes seizures is called **eclampsia**. When preeclampsia occurs during pregnancy, the babies may need to be delivered right away, even if they are not fully grown.

How can multiple pregnancy affect my risk of gestational diabetes?

Women carrying multiples have a high risk of **gestational diabetes**. This condition can increase the risk of preeclampsia and of developing **diabetes mellitus** later in life. Newborns may have breathing problems or low blood sugar levels. Diet, exercise, and sometimes medication can reduce the risk of these complications.

How can multiple pregnancy affect fetal growth?

Multiples are more likely to have growth problems than single babies. Multiples are called **discordant** if one fetus is much smaller than the others.

Discordant growth is common with multiples. It does not always signal a problem. Sometimes, though, a fetus's restricted growth may be caused by an infection, TTTS, or a problem with the placenta or umbilical cord. If growth restriction is suspected in one or both fetuses, frequent ultrasound exams may be done to track how the fetuses are growing.

Is screening for genetic disorders as accurate in multiple pregnancies?

Screening tests for **genetic disorders** that use a sample of a woman's blood are not as sensitive in multiple pregnancy. It is possible to have a positive screening test result when no problem is present in either fetus.

What diagnostic genetic tests are used during pregnancy?

Diagnostic tests for **birth defects** include **chorionic villus sampling (CVS)** and **amniocentesis**. These tests are harder to perform in multiples because each fetus

must be tested. There also is a small risk of loss of one or all of the fetuses. Results of these tests may show that one fetus has a disorder, while the others do not.

How can multiple pregnancy affect delivery?

The chance of cesarean birth is higher with multiples. In some cases, twins can be delivered by vaginal birth. How your babies are born depends on

- the number of babies and the position, weight, and health of each baby
- your health and how your labor is going
- the experience of your ob-gyn

Can multiple pregnancy affect my risk of postpartum depression?

Having multiples might increase your risk of [postpartum depression](#) . If you have intense feelings of sadness, anxiety, or despair that keep you from being able to do your daily tasks, talk with your ob-gyn or other member of your health care team.

Can I breastfeed if I have multiples?

Yes, but it may take some practice. Your milk supply will increase to the right amount. You will need to eat healthy foods and drink plenty of liquids. Lactation specialists are available at many hospitals and in your community to help you work out any problems you may have.

[\[How to Prepare for Breastfeeding in the Month Before Birth\]](#)

Glossary

Amniocentesis: A procedure in which amniotic fluid and cells are taken from the uterus for testing. The procedure uses a needle to withdraw fluid and cells from the sac that holds the fetus.

Amnionicity: The number of amniotic (inner) membranes that surround fetuses in a multiple pregnancy. When multiple fetuses have only one amnion, they share an amniotic sac.

Amniotic Sac: Fluid-filled sac in a woman's uterus. The fetus develops in this sac.

Birth Defects: Physical problems that are present at birth.

Cerebral Palsy: A disorder of the nervous system that affects movement, posture, and coordination. This disorder is present at birth.

Cesarean Birth: Birth of a fetus from the uterus through an incision made in the woman's abdomen.

Chorion: The outer membrane that surrounds the fetus.

Chorionic Villus Sampling (CVS): A procedure in which a small sample of cells is taken from the placenta and tested.

Chorionicity: The number of chorionic (outer) membranes that surround the fetuses in a multiple pregnancy.

Complication: A disease or condition that happens as a result of another disease or condition. An example is pneumonia that occurs as a result of the flu. A complication also can occur as a result of a condition, such as pregnancy. An example of a pregnancy complication is preterm labor.

Diabetes Mellitus: A condition in which the levels of sugar in the blood are too high.

Diagnostic Tests: Tests that look for a disease or cause of a disease.

Discordant: A large difference in the size of fetuses in a multiple pregnancy.

Eclampsia: Seizures occurring in pregnancy or after pregnancy that are linked to high blood pressure.

Egg: The female reproductive cell made in and released from the ovaries. Also called the ovum.

Embryo: The stage of development that starts at fertilization (joining of an egg and sperm) and lasts up to 8 weeks.

Fetus: The stage of human development beyond 8 completed weeks after fertilization.

Fraternal Twins: Twins that have developed from two different fertilized eggs.

Genetic Disorders: Disorders caused by a change in genes or chromosomes.

Gestational Diabetes: Diabetes that starts during pregnancy.

Identical Twins: Twins that have developed from a single fertilized egg that are usually genetically identical.

In Vitro Fertilization (IVF): A procedure in which an egg is removed from a woman's ovary, fertilized in a laboratory with the man's sperm, and then transferred to the woman's uterus to achieve a pregnancy.

Menstrual Cycle: The monthly process of changes that occur to prepare a woman's body for possible pregnancy. A menstrual cycle is defined as the first day of menstrual bleeding of one cycle to the first day of menstrual bleeding of the next cycle.

Multiple Pregnancy: A pregnancy where there are two or more fetuses.

Obstetrician–Gynecologist (Ob-Gyn): A doctor with special training and education in women's health.

Ovaries: Organs in women that contain the eggs necessary to get pregnant and make important hormones, such as estrogen, progesterone, and testosterone.

Ovulation: The time when an ovary releases an egg.

Placenta: An organ that provides nutrients to and takes waste away from the fetus.

Postpartum Depression: A type of depressive mood disorder that develops in the first year after the birth of a child. This type of depression can affect a woman's ability to take care of her child.

Preeclampsia: A disorder that can occur during pregnancy or after childbirth in which there is high blood pressure and other signs of organ injury. These signs include an abnormal amount of protein in the urine, a low number of platelets, abnormal kidney or

liver function, pain over the upper abdomen, fluid in the lungs, or a severe headache or changes in vision.

Prenatal Care: A program of care for a pregnant woman before the birth of her baby.

Preterm: Less than 37 weeks of pregnancy.

Screening Tests: Tests that look for possible signs of disease in people who do not have signs or symptoms.

Sperm: A cell made in the male testicles that can fertilize a female egg.

Twin–Twin Transfusion Syndrome (TTTS): A condition of identical twins in which one twin gets more blood than the other during pregnancy.

Ultrasound Exam: A test in which sound waves are used to examine inner parts of the body. During pregnancy, ultrasound can be used to check the fetus.

Umbilical Cord: A cord-like structure containing blood vessels. It connects the fetus to the placenta.

Uterus: A muscular organ in the female pelvis. During pregnancy, this organ holds and nourishes the fetus. Also called the womb.

If you have further questions, contact your ob-gyn.

Don't have an ob-gyn? [Learn how to find a doctor near you.](#)

FAQ188

Last updated: February 2021

Last reviewed: January 2023

Copyright 2024 by the American College of Obstetricians and Gynecologists. All rights reserved. Read [copyright and permissions information.](#)

This information is designed as an educational aid for the public. It offers current information and opinions related to women's health. It is not intended as a statement of the standard of care. It does not explain all of the proper treatments or methods of care. It is not a substitute for the advice of a physician. Read [ACOG's complete disclaimer](#).

About ACOG

Disclaimer

Contact Us

How to Find an Ob-Gyn



Copyright 2024 American College of Obstetricians and Gynecologists

Privacy Statement

|

Terms and Conditions of Use