

FAQs Nutrition During Pregnancy

Frequently Asked Questions

Overview

Why is nutrition during pregnancy important?

Eating well is one of the best things you can do during pregnancy. Good nutrition helps you handle the extra demands on your body as your pregnancy progresses. The goal is to balance getting enough nutrients to support the growth of your fetus and maintaining a healthy weight.

How much should I eat during pregnancy?

The popular saying is that you should "eat for two" when you're pregnant, but now we know that it's dangerous to eat twice your usual amount of food during pregnancy. Instead of "eating for two," think of it as eating twice as healthy.

If you are pregnant with one fetus, you need an extra 340 calories per day starting in the second trimester (and a bit more in the third trimester). That's roughly the calorie count of a glass of skim milk and half a sandwich.

If you are carrying twins, you should get about 600 extra calories a day. If you are carrying triplets, you should take in 900 extra calories a day.

Key Vitamins and Minerals

Why should I take a prenatal vitamin?

Vitamins and minerals play important roles in all of your body functions. Eating healthy foods and taking a prenatal vitamin every day should supply all the vitamins and minerals you need during pregnancy.

How many prenatal vitamins should I take each day?

Take only one serving of your prenatal supplement each day. Read the bottle to see how many pills make up one daily serving. If your obstetrician-gynecologist (ob-gyn) thinks you need an extra amount of a vitamin or mineral, your ob-gyn may recommend it as a separate supplement.

Can I take more prenatal vitamins to make up for a deficiency?

No, do not take more than the recommended amount of your prenatal vitamin per day. Some multivitamin ingredients, such as vitamin A, can cause birth defects at higher doses.

What vitamins and minerals do I need during pregnancy?

During pregnancy you need folic acid, iron, calcium, vitamin D, choline, omega-3 fatty acids, B vitamins, and vitamin C. See the below table for recommended amounts.

Key Vitamins and Minerals During Pregnancy

Nutrient (Daily Recommended Amount)	Why You and Your Fetus Need It	Best Sources
Calcium (1,300	Builds strong bones	Milk, cheese, yogurt, sardines, dark
milligrams for ages 14 to	and teeth	green leafy vegetables
18; 1,000 milligrams for		
ages 19 to 50)		
Iron (27 milligrams)	Helps red blood cells	Lean red meat, poultry, fish, dried
	deliver oxygen to your	beans and peas, iron-fortified
	fetus	cereals, prune juice
lodine (220 micrograms)	Essential for healthy	lodized table salt, dairy products,
· · · · · · · · · · · · · · · · · · ·	brain development	seafood, meat, some breads, eggs

Nutrient (Daily Recommended Amount)	Why You and Your Fetus Need It	Best Sources
Choline (450 milligrams)	•	Milk, beef liver, eggs, peanuts, soy products
Vitamin A (750 micrograms for ages 14 to 18; 770 micrograms for ages 19 to 50)	-	Carrots, green leafy vegetables, sweet potatoes
	Helps with bone growth	
Vitamin C (80 milligrams for ages 14 to 18; 85 milligrams for ages 19 to 50)	-	Citrus fruit, broccoli, tomatoes, strawberries
Vitamin D (600 international units)	Builds your fetus's bones and teeth	Sunlight, fortified milk, fatty fish such as salmon and sardines
	Helps promote healthy eyesight and skin	
Vitamin B6 (1.9 milligrams)	Helps form red blood cells	Beef, liver, pork, ham, whole-grain cereals, bananas
	Helps body use protein, fat, and carbohydrates	
Vitamin B12 (2.6 micrograms)	Maintains nervous system	Meat, fish, poultry, milk (vegetarians should take a supplement)

Nutrient (Daily Recommended Amount)	Why You and Your Fetus Need It	Best Sources
	Helps form red blood	
	cells	
Folic acid (600	Helps prevent birth	Fortified cereal, enriched bread and
micrograms)	defects of the brain	pasta, peanuts, dark green leafy
	and spine	vegetables, orange juice, beans.
		Also, take a daily prenatal vitamin
	Supports the general	with 400 micrograms of folic acid.
	growth and	
	development of the	
	fetus and placenta	

What is folic acid?

Folic acid, also known as folate, is a B vitamin that is important during pregnancy. Folic acid may help prevent major birth defects of the fetus's brain and spine called neural tube defects (NTDs).

How much folic acid should I take?

When you are pregnant you need 600 micrograms of folic acid each day. Because it's hard to get this much folic acid from food alone, you should take a daily prenatal vitamin with at least 400 micrograms starting at least 1 month before pregnancy and during the first 12 weeks of pregnancy.

If you have already had a child with an NTD, you should take 4 milligrams (mg) of folic acid each day as a separate supplement at least 3 months before pregnancy and for the first 3 months of pregnancy. You and your ob-gyn can discuss whether you need to supplement with more than 400 micrograms daily.

Why is iron important during pregnancy?

Iron is used by your body to make the extra blood that you and your fetus need during pregnancy. When you are not pregnant, you need 18 mg of iron per day. When you are pregnant, you need 27 mg per day. You can get this amount in most prenatal vitamins.

How can I make sure I'm getting enough iron?

In addition to taking a prenatal vitamin with iron, you should eat iron-rich foods such as beans, lentils, enriched breakfast cereals, beef, turkey, liver, and shrimp. You should also eat foods that help your body absorb iron, including oranges, grapefruit, strawberries, broccoli, and peppers.

Your blood should be tested during pregnancy to check for anemia. If you have anemia, your ob-gyn may recommend extra iron supplements.

What is calcium and what foods contain it?

Calcium is a mineral that builds your fetus's bones and teeth. If you are 18 or younger, you need 1,300 mg of calcium per day. If you are 19 or older, you need 1,000 mg per day.

Milk and other dairy products, such as cheese and yogurt, are the best sources of calcium. If you have trouble digesting milk products, you can get calcium from other sources, such as broccoli, fortified foods (cereals, breads, and juices), almonds and sesame seeds, sardines or anchovies with the bones, and dark green leafy vegetables. You can also get calcium from calcium supplements.

What is vitamin D and what foods contain it?

Vitamin D works with calcium to help the fetus's bones and teeth develop. Vitamin D is also essential for healthy skin and eyesight. Whether you are pregnant or not, you need 600 international units of vitamin D a day.

Good sources of vitamin D include fortified milk and breakfast cereal, fatty fish (salmon and mackerel), fish liver oils, and egg yolks.

How will I know if I'm getting enough vitamin D?

Many people do not get enough vitamin D. If your ob-gyn thinks you may have low levels of vitamin D, a test can be done to check the level in your blood. If it is below normal, you may need to take a vitamin D supplement.

What is choline and what foods contain it?

Choline plays a role in your fetus's brain development. It may also help prevent some common birth defects. Experts recommend that you get 450 mg of choline each day during pregnancy.

Choline can be found in chicken, beef, eggs, milk, soy products, and peanuts. Although the body produces some choline on its own, it doesn't make enough to meet all your needs while you are pregnant. It's important to get choline from your diet because it is not found in most prenatal vitamins.

What are omega-3 fatty acids and what foods contain omega-3s?

Omega-3 fatty acids are a type of fat found naturally in many kinds of fish. Omega-3s may be important for brain development before and after birth.

Flaxseed (ground or as oil) is also a good source of omega-3s. Other sources of omega-3s include broccoli, cantaloupe, kidney beans, spinach, cauliflower, and walnuts.

How much fish should I eat to get the omega-3 fatty acids that I need?

Eat two or three servings of fish or shellfish per week before getting pregnant, while pregnant, and while breastfeeding. A serving of fish is 4 ounces (oz).

Which types of fish should I avoid?

Some types of fish have higher levels of mercury than others. Mercury is a metal that has been linked to birth defects. Do not eat bigeye tuna, king mackerel, marlin, orange roughy, shark, swordfish, or tilefish. Limit white (albacore) tuna to only 6 oz a week. Also check advisories about fish caught in local waters.

What are B vitamins and what foods contain them?

B vitamins, including B1, B2, B6, B9, and B12, are key nutrients during pregnancy. These vitamins

- give you energy
- supply energy for your fetus's development
- promote good vision
- help build the placenta

Your prenatal vitamin should have the right amount of B vitamins that you need each day. Eating foods high in B vitamins is a good idea too, including liver, pork, chicken, bananas, beans, and whole-grain cereals and breads.

What is vitamin C and what foods contain it?

Vitamin C is important for a healthy immune system. It also helps build strong bones and muscles. During pregnancy, you should get at least 85 mg of vitamin C each day if you are older than 19, and 80 mg if you are younger than 19.

You can get the right amount of vitamin C in your daily prenatal vitamin, and also from citrus fruits and juices, strawberries, broccoli, and tomatoes.

How can I get enough water during pregnancy?

Drink throughout the day, not just when you are thirsty. Aim for 8 to 12 cups of water a day during pregnancy.

Planning Healthy Meals

How can I plan healthy meals during pregnancy?

There are many tools that can help you plan healthy meals. One useful tool is the MyPlate food-planning guide from the U.S. Department of Agriculture. The MyPlate website, www.myplate.gov, can help you learn how to make healthy food choices at every meal.

The MyPlate website offers a MyPlate Plan, which shows how much to eat based on how many calories you need each day. The MyPlate Plan is personalized based on your

- height
- prepregnancy weight
- physical activity level

The MyPlate Plan can help you learn about choosing foods from each food group to get the vitamins and minerals you need during pregnancy. The MyPlate Plan can also help you limit calories from added sugars and saturated fats.

What are the five food groups?

- Grains
- Fruits
- Vegetables
- Protein foods
- Dairy foods

What are grains?

Bread, pasta, oatmeal, cereal, and tortillas are all grains. Whole grains are those that haven't been processed and include the whole grain kernel. Oats, barley, quinoa, brown rice, and bulgur are all whole grains, as are products made with those grains. Look for the words "whole grain" on the product label. When you plan meals, make half of your grain servings whole grains.

What types of fruit should I eat?

You can eat fresh, canned, frozen, or dried fruit. Juice that is 100 percent fruit juice counts in the fruit category, but it is best to eat mostly whole fruit instead of juice. Make half your plate fruit and vegetables during mealtimes.

What types of vegetables should I eat?

You can eat raw, canned, frozen, or dried vegetables or drink 100 percent vegetable juice. Use dark leafy greens to make salads. Make half your plate fruit and vegetables during mealtimes.

What are protein foods?

Meat, poultry, seafood, beans, peas, eggs, processed soy products, nuts, and seeds all contain protein. Eat a variety of protein each day.

What are dairy foods?

Milk and milk products, such as cheese and yogurt, make up the dairy group. Make sure any dairy foods you eat are pasteurized. Choose fat-free or low-fat (1 percent) varieties.

Why are oils and fats important?

Oils and fats are another part of healthy eating. Although they are not a food group, they do give you important nutrients. During pregnancy, the fats that you eat provide energy and help build the placenta and many fetal organs.

What are healthy sources of oils and fats?

Oils in food come mainly from plant sources, such as olive oil, nut oils, and grapeseed oil. They can also be found in certain foods, such as some fish, avocados, nuts, and olives.

Most of the fats and oils in your diet should come from plant sources. Limit solid fats, such as those from animal sources. Solid fats can also be found in processed foods.

Weight Gain

How much weight should I gain during pregnancy?

Weight gain depends on your health and your body mass index (BMI) before you were pregnant. If you were underweight before pregnancy, you should gain more weight than those who had a normal weight before pregnancy. If you were overweight or obese before pregnancy, you should gain less weight. The amount of weight gain differs by trimester :

- During your first 12 weeks of pregnancy—the first trimester—you might gain only 1 to 5 pounds or none at all.
- If you were a healthy weight before pregnancy, you should gain a half-pound to 1 pound per week in your second and third trimesters.

See the below table for recommended weight gain during pregnancy.

Body Mass Index (BMI) Before Pregnancy	Rate of Weight Gain in the Second and Third Trimesters* (Pounds Per Week)	Recommended Total Weight Gain With a Single Fetus (in Pounds)	Recommended Total Weight Gain With Twins (in Pounds)
Less than 18.5 (underweight)	1.0 to 1.3	28 to 40	Not known
18.5 to 24.9 (normal weight)	0.8 to 1.0	25 to 35	37 to 54
25.0 to 29.9 (overweight)	0.5 to 0.7	15 to 25	31 to 50
30.0 and above (obese)	0.4 to 0.6	11 to 20	25 to 42

*Assumes a first-trimester weight gain between 1.1 and 4.4 pounds

Source: Institute of Medicine and National Research Council. 2009. Weight Gain During Pregnancy: Reexamining the Guidelines. Washington, DC: The National Academies Press.

How many extra calories should I eat?

During the first trimester with one fetus, usually no extra calories are needed. In the second trimester, you will need an extra 340 calories per day, and in the third trimester, about 450 extra calories a day. To get the extra calories during the day, have healthy snacks on hand, such as nuts, yogurt, and fresh fruit.

How can being overweight cause problems during pregnancy?

Excess weight during pregnancy is associated with several pregnancy and childbirth complications, including

- high blood pressure
- preeclampsia
- preterm birth
- gestational diabetes

Obesity during pregnancy also increases the risk of:

• a larger than normal fetus (macrosomia)

- birth injury
- cesarean birth
- birth defects, especially NTDs

What if I am overweight?

You and your ob-gyn should work together to develop a nutrition and exercise plan. If you are gaining less than what the guidelines suggest, and if your fetus is growing well, gaining less than the recommended guidelines can have benefits. If your fetus is not growing well, changes may need to be made to your diet and exercise plan.

Resources and Glossary

Resources

MyPlate

Healthy eating resources from the U.S. Department of Agriculture (USDA).

www.myplate.gov

- Healthy Eating on a Budget: https://www.myplate.gov/eat-healthy/healthy-eating-budget
- Pregnancy and Breastfeeding: https://www.myplate.gov/life-stages/pregnancy-and-breastfeeding
- MyPlate Plan: https://www.myplate.gov/myplate-plan

Food Sources of Select Nutrients

Examples of foods that are good sources of important nutrients. www.dietaryguidelines.gov/resources/2020-2025-dietary-guidelines-onlinematerials/food-sources-select-nutrients

Glossary

Anemia: Abnormally low levels of red blood cells in the bloodstream. Most cases are caused by iron deficiency (lack of iron).

Birth Defects: Physical problems that are present at birth.

Body Mass Index: A number calculated from height and weight. BMI is used to determine whether a person is underweight, normal weight, overweight, or obese.

Calories: Units of heat used to express the fuel or energy value of food.

Cesarean Birth: Birth of a fetus from the uterus through an incision (cut) made in the woman's abdomen.

Complications: Diseases or conditions that happen as a result of another disease or condition. An example is pneumonia that occurs as a result of the flu. A complication also can occur as a result of a condition, such as pregnancy. An example of a pregnancy complication is preterm labor.

Fetus: The stage of human development beyond 8 completed weeks after fertilization.

Folic Acid: A vitamin that reduces the risk of certain birth defects when taken before and during pregnancy.

Gestational Diabetes: Diabetes that starts during pregnancy.

High Blood Pressure: Blood pressure above the normal level. Also called hypertension.

Macrosomia: A condition in which a fetus grows more than expected, often weighing more than 8 pounds and 13 ounces (4,000 grams).

Neural Tube Defects: Birth defects that result from a problem in development of the brain, spinal cord, or their coverings.

Obstetrician–Gynecologist (Ob-Gyn): A doctor with special training and education in women's health.

Oxygen: An element that we breathe in to sustain life.

Placenta: An organ that provides nutrients to and takes waste away from the fetus.

Preeclampsia: A disorder that can occur during pregnancy or after childbirth in which there is high blood pressure and other signs of organ injury. These signs include an abnormal amount of protein in the urine, a low number of platelets, abnormal kidney or

liver function, pain over the upper abdomen, fluid in the lungs, or a severe headache or changes in vision.

Preterm: Less than 37 weeks of pregnancy.

Trimester: A 3-month time in pregnancy. It can be first, second, or third.

If you have further questions, contact your ob-gyn.

Don't have an ob-gyn? Learn how to find a doctor near you.

FAQ001 Last updated: June 2023 Last reviewed: December 2021

Copyright 2024 by the American College of Obstetricians and Gynecologists. All rights reserved. Read copyright and permissions information.

This information is designed as an educational aid for the public. It offers current information and opinions related to women's health. It is not intended as a statement of the standard of care. It does not explain all of the proper treatments or methods of care. It is not a substitute for the advice of a physician. Read ACOG's complete disclaimer.

About ACOG

Disclaimer

Contact Us

How to Find an Ob-Gyn

f in 💿

Privacy Statement

Terms and Conditions of Use